



E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2018; SPI: 2481-2486

SS Porte
Department of Soil Science and
Agricultural Chemistry,
Jawaharlal Nehru Krishi
Vishwavidyalaya, Jabalpur,
(M.P.), India

B Sachidanand
Department of Soil Science and
Agricultural Chemistry,
Jawaharlal Nehru Krishi
Vishwavidyalaya, Jabalpur,
(M.P.), India

HK Rai
Department of Soil Science and
Agricultural Chemistry,
Jawaharlal Nehru Krishi
Vishwavidyalaya, Jabalpur,
(M.P.), India

Shyamal
Department of Soil Science and
Agricultural Chemistry,
Jawaharlal Nehru Krishi
Vishwavidyalaya, Jabalpur,
(M.P.), India

Vaishali Sharma
Department of Soil Science and
Agricultural Chemistry,
Jawaharlal Nehru Krishi
Vishwavidyalaya, Jabalpur,
(M.P.) India

Tarun Suryavanshi
Department of Soil Science and
Agricultural Chemistry,
Jawaharlal Nehru Krishi
Vishwavidyalaya, Jabalpur,
(M.P.), India

Kritika Dongre
Department of Soil Science and
Agricultural Chemistry,
Jawaharlal Nehru Krishi
Vishwavidyalaya, Jabalpur,
(M.P.), India

Correspondence

SS Porte
Department of Soil Science and
Agricultural Chemistry,
Jawaharlal Nehru Krishi
Vishwavidyalaya, Jabalpur,
(M.P.), India

Effects of targeted yield based fertilizer application on soil properties, growth, yield and quality of rice under rice-wheat cropping system in a Vertisol

SS Porte, B Sachidanand, HK Rai, Shyamal, Vaishali Sharma, Tarun Suryavanshi and Kritika Dongre

Abstract

The present investigation was carried out at the research farm of JNKVV under AICRP on STCR, Department of Soil Science and Agricultural Chemistry, College of Agriculture, JNKVV, Jabalpur (MP). The experiment was undertaken on Typic Hapluster soil on the field of soil science Department JNKVV Research Farm, Jabalpur. The experiment was taken under the ongoing project of AICRP on STCR, taking six different treatments based on targeted yield of rice and wheat cropping sequence. During *kharif* season in rice crop with 6 treatments as RT₁ (Control), RT₂ (GRD @ 80:50:30 NPK kg ha⁻¹), RT₃ (Targeted Yield 5 t ha⁻¹), RT₄ (Targeted Yield, 6 t ha⁻¹), RT₅ (Targeted Yield, 5 t ha⁻¹ with 5 t FYM ha⁻¹) and RT₆ (Targeted Yield 6 t ha⁻¹ with 5 t FYM ha⁻¹). During *rabi* season in wheat crop, 6 treatments as WT₁ (Control), WT₂ (GRD @ 120:80:60 NPK kg ha⁻¹), WT₃ (Targeted Yield 4.5 t ha⁻¹), WT₄ (Targeted Yield, 6 t ha⁻¹), WT₅ (Targeted Yield, 4.5 t ha⁻¹ with 5 t FYM ha⁻¹) and WT₆ (Targeted Yield 6 t ha⁻¹ with 5 t FYM ha⁻¹) were tested under randomized block design with four replications. The pH value recorded at 0-20 cm soil depth in rice crop at 30, 60, 90 DAS and at harvest under different treatments was slightly decreased than the initial soil pH value of 7.78. In case of wheat, different treatments did not give significant influence on pH of surface (0-20 cm) soils at different time interval. In case of rice, EC values at 0-20 cm depth showed increasing trend over passage of time of growth as compared to initial value. Wheat crop showed increasing trend of EC at 30 and 60 DAS and then decreasing trend there after again. Application of nutrient in integrated manner increased organic carbon at 0-20 cm at different time intervals in both rice and wheat. Maximum plant height, tillers/hills and penicle length were recorded under RT₆. The highest yield of grain and straw was recorded in treatment RT₆ having higher target yield 5.57 t ha⁻¹ and 8.53 t ha⁻¹ respectively. Maximum protein content was also recorded under RT₆.

Keywords: Targeted yield, soil properties, rice growth, yield attributes, yield and quality

1. Introduction

Rice (*Oryza sativa* L.) is one of the most staple food crops, which supplies major source of calories for about 45% of world population particularly to the people of Asian countries. Rice stands second in the world after wheat in area and consumes 90% of world rice (Anonymous, 2006). Intensive agriculture for higher yields using synthetic fertilizer especially nitrogen and phosphorus enhances the production, but on the other hand addition of these fertilizers adversely effects the environment due to emission of greenhouse gases. Rice agriculture is the most important anthropogenic sources of methane. Soil fertility is a prerequisite to its productivity.

Quantity of fertilizer depends on uptake of nutrients by crops. It should be based on soil analysis. Dumping of fertilizers by the farmers in fields without information on soil fertility status and nutrient requirements by crops might cause adverse effects on soils and crops both regarding nutrient toxicity and deficiency either by over use or inadequate use. INM approach is flexible and minimizes use of chemicals but maximize use efficiency and improve the soil health. The targeted yield concept was developed by Ramamoorthy *et al.* (1967). Balanced NPK fertilization has received considerable attention in India (Gosh *et al.*, 2004; Hegde *et al.*, 2004 and Prasad *et al.*, 2004). It provides the balanced nutrition to the crop according to the actual requirement of the crop and soil fertility conditions. The soil test crop response (STCR) approach for targeted yield is unique in indicating both soil test based fertilizer dose and the level of yield that can be achieved with good agricultural practices.

Degradation of soil health has also been reported due to long-term imbalanced use of fertilizer nutrients. Although, overall nutrient use (N: P₂O₅:K₂O) of 4:2:1 is considered ideal for Indian soils, the present use ratio of 6.8:2.8:1 is far off the mark. This imbalance nutrient use has resulted in wide gap between crop removal and fertilizer application. Long- term experiments,

in India have in general showed that P and K status in soils at all centers has gone down when only N was applied. The partial factor productivity of fertilizers during the last three and half decades showed a declining trend from 48 kg food grains/kg NPK fertilizer in 1970-71 to 10 kg food grains/ kg NPK fertilizer in 2007-08 (Aulakh and Benbi, 2008, Subba Rao and Reddy, 2009). The soil test based fertilizer application is on the basis of nutrient required by the crop to produce substantial yield. Change in cropping sequence with respect to availability of resources the integrated approach of nutrient supply through inorganic and organic (FYM) has become very much promising in building soil health and quality of produce.

Materials and methods

The field experiment was conducted on rice crops during Kharif 2013 at Research Farm of Department of Soil Science & Agricultural Chemistry, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur (23°10' N latitude and 79°57' E longitude). The experimental soil was medium black belonging to fine montmorillonitic hyperthermic family of Typic Haplustert and had pH of 7.8, electrical conductivity 0.25 dSm⁻¹ (1: 2.5 soil: water ratio) and organic carbon content 0.55%. Treatments schedule for rice crop of RT₁-Control, RT₂-General Recommended Dose (GRD), RT₃ - Targeted Yield 5.0 t ha⁻¹ RT₄- Targeted Yield 6.0 t ha⁻¹ RT₅- Targeted Yield 5.0 t ha⁻¹ + 5 t FYM ha⁻¹ and RT₆- Targeted Yield 6.0 t ha⁻¹ + FYM 5 t ha⁻¹. During *rabi* season in wheat crop, 6 treatments as WT₁ (Control), WT₂ (GRD @ 120:80:60 NPK kg ha⁻¹), WT₃ (Targeted Yield 4.5 t ha⁻¹), WT₄ (Targeted

Yield, 6 t ha⁻¹), WT₅ (Targeted Yield, 4.5 t ha⁻¹ with 5 t FYM ha⁻¹) and WT₆ (Targeted Yield 6 t ha⁻¹ with 5 t FYM ha⁻¹) There were six treatments replicated four times in a randomized block design for the crop. Treated seeds of rice (Kranti) were sown in rows at proper spacing in the first week of July 2013, after basal application of fertilizers as per treatments. The soil samples were collected before sowing and after the harvest of rice crop during 2013 with the help of a tube auger (stainless steel) from each plot at 0-20 cm soil depth. Basic soil parameters were estimated by using standard laboratory procedures (Jackson, 1973).

To compute fertilizer doses for any yield target based on soil test value fertilizer adjustment equations were used (Table 1 and 2) as per procedure of Ramamoorthy *et al.* 1967. The targeted yield for rice was 5.0 and 6.0 tones ha⁻¹ and was fixed as 80:50:30 kg ha⁻¹ with NPK dose. The fertilizer materials used were FYM, urea, single super phosphate and Murat of potash. Full dose of P and K and half dose of N were applied and mixed thoroughly with soil at the time of sowing. The remaining half dose of N was top-dressed in two splits at tillering stage and boot stage. Cultivation of rice was done adopting proper package of practices. All climatic conditions were favourable for growth and development of crop. The rice crop was grown under irrigated soil condition. The grain yields of rice were recorded at the harvest of crop on maturity for each treatment. The soil and plant samples were analyzed by standard laboratory procedure. Thus the analysis of variance was carried out using the randomized block design as described by Gomez and Gomez, 1984.

Table 1: Soil test value for rice-wheat cropping sequence (as per table)

S.N.	Nutrients	Soil test value (kg ha ⁻¹)			
		Rice	Wheat	Rice	Wheat
		2013	2013-14	2014	2014-15
1.	Available soil N (kg ha ⁻¹)	230.04	196.55	223.25	195.02
2.	Available soil P (kg ha ⁻¹)	17.06	18.35	17.59	16.43
3.	Available soil K (kg ha ⁻¹)	250.74	226.34	250.58	230.46

Table 2: Fertilizer adjustment equation used for rice and wheat

a. Rice					b. Wheat				
F N	=	4.25 T	-	0.45 SN	F N	=	4.40 T	-	0.45 SN
F P ₂ O ₅	=	3.55 T	-	4.89 SP	F P ₂ O ₅	=	4.00 T	-	5.73 SP
F K ₂ O	=	2.10 T	-	0.18 SK	F K ₂ O	=	2.53 T	-	0.16 SK

Where

F N= Fertilizer nitrogen (kg ha-1)

F P₂O₅=Fertilizer phosphorus (kg ha-1)

F K₂O=Fertilizer potassium (kg ha-1)

T= Desired yield target (q ha-1)

SN= Available soil nitrogen (kg ha-1)

SP= Available soil phosphorus (kg ha-1)

SK= Available soil potassium (kg ha-1)

Table 3: Fertilizer requirements for rice-wheat cropping sequence (2013-14)

Treatment code	Treatment details	Nutrient supplied (kg ha ⁻¹)			FYM (t ha ⁻¹)
		N	P ₂ O ₅	K ₂ O	
(a) Kharif (Rice)					
RT ₁	Control	0	0	0	0
RT ₂	General recommended dose	80	50	30	0
RT ₃	Targeted Yield 5 t ha ⁻¹	119.12	100.73	68.85	0
RT ₄	Targeted Yield 6 t ha ⁻¹	161.62	136.23	89.85	0
RT ₅	Targeted yield 5 t ha ⁻¹ + 5 t FYM ha ⁻¹	69.12	50.73	18.85	5
RT ₆	Targeted Yield 6 t ha ⁻¹ + 5 t FYM ha ⁻¹	111.62	86.13	32.85	5
(b) Rabi (Wheat)					
WT ₁	Control	0	0	0	0
WT ₂	General recommended dose	120	80	60	0
WT ₃	Targeted Yield 4.5 t ha ⁻¹	117.72	82.99	78.90	0
WT ₄	Targeted Yield 6 t ha ⁻¹	178.61	145.57	119.54	0
WT ₅	Targeted Yield 4.5 t ha ⁻¹ + 5 t FYM ha ⁻¹	107.68	89.69	79.08	5
WT ₆	Targeted Yield 6 t ha ⁻¹ + 5 t FYM ha ⁻¹	183.72	145.05	115.60	5

Table 4: Fertilizer requirements for rice-wheat cropping sequence (2014-15)

Treatment code	Treatment details	Nutrient supplied (kg ha ⁻¹)			FYM (t ha ⁻¹)
		N	P ₂ O ₅	K ₂ O	
(a) Kharif (Rice)					
R T ₁	Control	0	0	0	0
R T ₂	General recommended dose (GRD)	80	50	30	0
R T ₃	Targeted Yield 5 t ha ⁻¹	82.67	99.156	81.07	0
R T ₄	Targeted Yield 6 t ha ⁻¹	111.05	141.753	96.54	0
R T ₅	Targeted yield 5 t ha ⁻¹ + FYM 5 t ha ⁻¹	88.31	92.463	73.16	5
R T ₆	Targeted Yield 6 t ha ⁻¹ + FYM 5 t ha ⁻¹	122.34	153.244	90.09	5
(b) Rabi (Wheat)					
W T ₁	Control	0	0	0	0
W T ₂	General recommended dose (GRD)	120	80	60	0
W T ₃	Targeted Yield 4.5 t ha ⁻¹	130.26	100.76	91.99	0
W T ₄	Targeted Yield 6 t ha ⁻¹	193.75	142.76	131.01	0
W T ₅	Targeted Yield 4.5 t ha ⁻¹ + FYM 5 t ha ⁻¹	122.73	55.03	91.63	5
W T ₆	Targeted Yield 6 t ha ⁻¹ + FYM 5 t ha ⁻¹	206.30	149.69	130.83	5

Results and Discussion

Soil properties

Soil pH

Soil pH is an intrinsic property which is decided by the exchangeable cations on clay surface and taken larger time to get change. The data on pH of surface (0-20 cm) soil recorded at 30, 60, 90 DAS at harvest of rice and wheat are presented in (Table 5). In general, the pH value recorded at 0-20 cm soil in rice crop at 30, 60, 90 DAS and at harvest under different treatments was slightly decreased than the initial soil pH value of 7.78 during both the years and on mean basis. The data recorded on pH at 0-20 cm soil of rice crop revealed that different treatments did not have significant influence at different interval during both the years and on mean basis. However, maximum pH was noted under control (RT₁) followed by GRD @ 80:50:30 NPK kg ha⁻¹ (RT₂) during both the years and on mean basis. Other treatments had almost similar values of pH at 0-20 cm (pH 7.65 to 7.69) at different interval during both the years and on mean basis.

In case of wheat (Table 6), data revealed that different treatments did exert significant influence on pH in surface (0-20 cm) soils at different time interval during both the years and on mean basis. However, maximum pH was noted in control (WT₁) at different interval during both the years and on mean basis. At 90 DAS and at harvest stage, treatments T. Y. 4.5 t + 5 t FYM ha⁻¹ (WT₅) and T. Y. 6.0 t + 5 t FYM ha⁻¹ (WT₆) recorded relatively lower pH as compared to others during both the years and on mean basis. In general, the pH at 0-20 cm soil in (wheat) at 30, 60, 90 DAS and at harvest in different treatments significantly decreased than the initial pH value of 7.77 during both the years and on mean basis

Electrical conductivity (dSm⁻¹)

The data on EC of surface (0-20 cm depth) soils recorded at 30, 60, 90 DAS and at harvest of rice and wheat are presented in (Table 7). In general, the EC at 0-20 cm soil depth in rice at 30, 60, 90 DAS and at harvest under different treatments was increased as compared to the initial soil EC value of 0.22 (dSm⁻¹) during both the years and on mean basis. Further, EC values in soil at 0-20 cm revealed slightly increasing trend from 30 DAS to at harvest. The data on EC at 0-20 cm soil depth in rice crop revealed that different treatments did not exert significant influence on this parameter at different stages of rice during both the years and on mean basis. However, at 0-20 cm depth of soil at harvest stage, numerically maximum EC value was noted under T.Y. 5.0 t ha⁻¹ + 5 t FYM ha⁻¹ (RT₅) and T.Y.6.0 t ha⁻¹ (RT₄) on mean basis.

As regards to wheat crop, different treatments failed to have

significant impact on EC values at 0-20 cm of soil during both the years and on mean basis. In general, at 0-20 cm soil, the EC values increased from 30 DAS to 60 DAS, thereafter it declined at 90 DAS and again it increased slightly at harvest during both the years and on mean basis. The maximum EC values at 0-20 cm soil depth were noted at 60 DAS in all the treatments during both the years and on mean basis. At harvest stage, numerically maximum EC value was noted in T. Y. 6.0 t ha⁻¹ (WT₄) followed by T. Y. 4.5 t ha⁻¹ + 5 t FYM ha⁻¹ (WT₅), T. Y. 4.5 t ha⁻¹ (WT₃) and T. Y. 6.0 t ha⁻¹ + 5 t FYM ha⁻¹ (WT₆) on mean basis at 0-20 cm soil depth (Table 8).

Soil organic carbon (g kg⁻¹)

Soil organic carbon is key soil property for evaluating the soil health. The data on soils organic carbon of surface (0-20 cm) soil at 30, 60, 90 DAS and at harvest of rice and wheat are presented in (Table 9).

The data on organic carbon in soil at 0-20 cm revealed that different treatments had significant effect on this parameter at different time interval of rice during both the years and on mean basis. Treatment T.Y.6.0 t ha⁻¹ + 5 t FYM ha⁻¹ (RT₆) gave significantly higher organic carbon at 0-20 cm depth at different time interval as compared to others during both the years and on mean basis. However, at 60 DAS, it was at par to treatment T.Y.5.0 t ha⁻¹ + 5 t FYM ha⁻¹ (RT₅) during 2014 and on mean basis (Table 10). At 20-40 cm soil at various time interval of rice, treatment T.Y.6.0 t ha⁻¹ + 5 t FYM ha⁻¹ (RT₆) showed significantly higher organic carbon as compared to others during both the years and on mean basis. However, at harvest during 2013, it was comparable to treatment T.Y.5.0 t ha⁻¹ + 5 t FYM ha⁻¹ (RT₅).

As regards to wheat crop, the organic carbon at 0-20 cm depth of soil had significantly higher value in treatment T.Y.6.0 t ha⁻¹ + 5 t FYM ha⁻¹ (WT₆) as compared to others at different time interval during both the years and on mean basis. However, it was at par to treatment T.Y.4.5 t ha⁻¹ + 5 t FYM ha⁻¹ (WT₅) at 30 DAS during 2013 and on mean basis and at 60 DAS during 2014. Further, at 30 DAS, treatments T.Y.6.0 t ha⁻¹ (WT₄) and T.Y.4.5 t ha⁻¹ (WT₃) also recorded at par values of organic carbon on mean basis only (Table 10).

Growth parameters

The data presented in Table 11 shows that plant height was maximum in RT₆ whereas minimum plant height was recorded in RT₁ (44.85 cm) followed by RT₂ (63.95 cm). However, RT₃, RT₄ and RT₅ were at par with RT₆ in both year of cropping sequence (2013 and 2014). Similar result was also

reported by Sahu *et al.*, (2015).

The data recorded on number of tillers hill⁻¹ showed significant variation. It is obvious from the data presented in table 5 that number of tillers hill⁻¹ was found maximum (9.29) under RT₆ which was at par with RT₃, RT₄ and RT₅ whereas, the minimum number of tillers hill⁻¹ (4.35) was recorded in RT₁ followed by RT₂ respectively in both year of cropping sequence. Similar result was also reported by Tabar *et al.*, (2012).

Data on panicles hill⁻¹ of rice as affected by different treatments are given in Table 11. The RT₆ had significantly higher number of tillers and panicle hill⁻¹ (9.23) over all the treatments except RT₃, RT₄ and RT₅ which was being at par in both years of cropping sequence. However, the minimum number of tillers and panicle hill⁻¹ (4.15) associated with RT₁ (control) followed by (RT₂), respectively in both year of cropping sequence. Similar results were also reported by Sahu *et al.* (2015) and Chaubey *et al.*, (2015).

Data on panicle length of rice as affected by different treatments are presented in Table 11. Data on panicles hill⁻¹ of rice as affected by different treatments are given in Table 11. The RT₆ had significantly higher number of tillers and panicle hill⁻¹ (9.23) over all the treatments except RT₃, RT₄ and RT₅ which was being at par in both years of cropping sequence. However, the minimum number of tillers and panicle hill⁻¹ (4.15) associated with RT₁ (control) followed by (RT₂), respectively in both year of cropping sequence. Similar results were also reported by Tan *et al.*, (2000).

Grain and straw yields

A perusal of data of grain and straw yield of paddy given in

Table 12 indicated significant variation due to different treatments and were found significantly higher over control. Higher target of 6 t ha⁻¹ (T.Y.6 t ha⁻¹ + 5 t ha⁻¹ FYM) could not be achieved and deviated by ± 6.17% negatively, whereas, the target of 5 t ha⁻¹ (T.Y.5 t ha⁻¹ + 5 t ha⁻¹ FYM) was obtained comfortably. The grain yield target was achieved only in treatment RT₅ (T.Y.5 t ha⁻¹ + 5 t ha⁻¹ FYM) which was significantly increased over control having increased by 60.32 percent. The treatment RT₆ target (T.Y.6 t ha⁻¹ + 5 t ha⁻¹ FYM) could not be achieved however; it resulted significantly superior over rest of the treatments except RT₄ and RT₅ in both years of cropping sequence. The yield increased over GRD due to 6 t ha⁻¹ target (T.Y.6 t ha⁻¹ + 5 t ha⁻¹ FYM) was 25.04 percent. This result confirms with the result reported by Roy *et al.*, (1997), Mishra and Vyas (1992). The overall increase in yield due to treatments either GRD or soil test based fertilizer fix application for fix target with and without FYM have markedly augmented yield of paddy. Pandya *et al.* (2005), Dwivedi *et al.*, (2011), Mishra *et al.*, (2015) and Sahu *et al.*, (2015).

Protein content

It was observed that the protein content gradually increased ranging from 5.97 to 7.70 percent differing significantly with control. The highest protein content was recorded in RT₆ (TY 6 t ha⁻¹ + 5 t ha⁻¹ FYM) in both year of cropping sequence. Similar finding have been also reported by Tayefe *et al.*, (2012) (Table 12).

Table 5: pH of rice- growing soil (0-20cm) at different time interval as influenced by targeted yield- based nutrient application

Treatment	pH at 0-20 cm of rice- growing soil											
	30 DAS			60 DAS			90 DAS			At harvest		
	2013	2014	Mean	2013	2014	Mean	2013	2014	Mean	2013	2014	Mean
RT ₁ - Control	7.69	7.69	7.69	7.69	7.69	7.69	7.69	7.68	7.69	7.69	7.68	7.69
RT ₂ - GRD@ 80:50:30 NPK kg ha ⁻¹	7.68	7.68	7.68	7.69	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.67
RT ₃ - T.Y.5.0 t ha ⁻¹	7.67	7.67	7.67	7.68	7.68	7.68	7.68	7.67	7.68	7.68	7.67	7.67
RT ₄ - T.Y.6.0 t ha ⁻¹	7.67	7.67	7.67	7.68	7.68	7.68	7.68	7.67	7.68	7.68	7.67	7.67
RT ₅ - T.Y.5.0 t ha ⁻¹ +5 t FYM ha ⁻¹	7.67	7.66	7.67	7.67	7.67	7.67	7.67	7.67	7.67	7.67	7.67	7.66
RT ₆ - T.Y.6.0 t ha ⁻¹ + 5 t FYM ha ⁻¹	7.67	7.65	7.66	7.67	7.67	7.67	7.66	7.66	7.66	7.66	7.66	7.66
S Em ±	0.01	0.01	0.01	0.008	0.006	0.006	0.008	0.01	0.007	0.008	0.013	0.007
CD (p = 0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Table 6: pH of wheat- growing soil (0-20 cm) at different time interval as influenced by targeted yield-based nutrient application

Treatment	pH at 0-20 cm of wheat - growing soil											
	30 DAS			60 DAS			90 DAS			At harvest		
	2013-14	2014-15	Mean	2013-14	2014-15	Mean	2013-14	2014-15	Mean	2013-14	2014-15	Mean
WT ₁ - Control	7.69	7.69	7.69	7.69	7.69	7.69	7.69	7.69	7.69	7.69	7.69	7.69
WT ₂ - GRD@ 120:80:60 NPK kg ha ⁻¹	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68
WT ₃ - T.Y.4.5 t ha ⁻¹	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68	7.68
WT ₄ - T.Y.6.0 t ha ⁻¹	7.68	7.68	7.68	7.68	7.68	7.68	7.67	7.68	7.68	7.67	7.67	7.67
WT ₅ - T.Y.4.5 t ha ⁻¹ + 5 t FYM ha ⁻¹	7.68	7.68	7.68	7.68	7.67	7.67	7.66	7.66	7.66	7.67	7.66	7.66
WT ₆ - T.Y.6.0 t ha ⁻¹ + 5 t FYM ha ⁻¹	7.68	7.67	7.68	7.67	7.67	7.67	7.66	7.66	7.66	7.67	7.66	7.66
S Em ±	0.007	0.008	0.005	0.008	0.01	0.01	0.010	0.007	0.005	0.005	0.005	0.003
CD (P=0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Table 7: EC (dSm⁻¹) of rice- growing soil (0-20cm) at different time interval as influenced by targeted yield- based nutrient application

Treatment	EC (dSm ⁻¹) at 0-20 cm of rice - growing soil											
	30 DAS			60 DAS			90 DAS			At harvest		
	2013	2014	Mean	2013	2014	Mean	2013	2014	Mean	2013	2014	Mean
RT ₁ - Control	0.203	0.195	0.199	0.205	0.196	0.200	0.202	0.222	0.212	0.250	0.253	0.251
RT ₂ - GRD@ 80:50:30 NPK kg ha ⁻¹	0.230	0.218	0.224	0.231	0.220	0.225	0.206	0.230	0.218	0.243	0.240	0.241
RT ₃ - T.Y.5.0 t ha ⁻¹	0.218	0.205	0.211	0.222	0.208	0.215	0.207	0.233	0.220	0.245	0.258	0.251
RT ₄ - T.Y.6.0 t ha ⁻¹	0.233	0.218	0.225	0.233	0.220	0.226	0.220	0.225	0.223	0.248	0.258	0.253
RT ₅ - T.Y.5.0 t ha ⁻¹ +5 t FYM ha ⁻¹	0.218	0.198	0.208	0.223	0.203	0.213	0.207	0.222	0.214	0.235	0.270	0.253
RT ₆ - T.Y.6.0 t ha ⁻¹ + 5 t FYM ha ⁻¹	0.228	0.203	0.215	0.226	0.211	0.218	0.214	0.223	0.218	0.235	0.258	0.246
S Em ±	0.008	0.006	0.005	0.006	0.006	0.004	0.006	0.010	0.007	0.006	0.005	0.004
CD (p = 0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Table 8: EC (dSm⁻¹) of wheat - growing soil (0-20cm) at different time interval as influenced by targeted yield - based nutrient application

Treatment	EC (dSm ⁻¹) at 0-20 cm of wheat - growing soil											
	30 DAS			60 DAS			90 DAS			At harvest		
	2013-14	2014-14	Mean	2013-14	2014-15	Mean	2013-14	2014-15	Mean	2013-14	2014-15	Mean
WT ₁ - Control	0.190	0.193	0.191	0.249	0.249	0.249	0.211	0.215	0.213	0.228	0.234	0.231
WT ₂ - GRD@ 120:80:60 NPK kg ha ⁻¹	0.200	0.205	0.203	0.253	0.253	0.253	0.205	0.210	0.207	0.233	0.238	0.235
WT ₃ - T.Y.4.5 t ha ⁻¹	0.198	0.208	0.203	0.259	0.258	0.259	0.210	0.217	0.213	0.233	0.240	0.236
WT ₄ - T.Y.6.0 t ha ⁻¹	0.195	0.210	0.203	0.261	0.261	0.261	0.211	0.216	0.213	0.238	0.242	0.240
WT ₅ - T.Y.4.5 t ha ⁻¹ + 5 t FYM ha ⁻¹	0.218	0.210	0.214	0.262	0.264	0.263	0.204	0.212	0.208	0.230	0.243	0.237
WT ₆ - T.Y.6.0 t ha ⁻¹ + 5 t FYM ha ⁻¹	0.210	0.208	0.209	0.263	0.265	0.264	0.210	0.216	0.213	0.235	0.238	0.236
S Em ±	0.006	0.006	0.004	0.002	0.002	0.001	0.012	0.008	0.009	0.004	0.004	0.003
CD (p = 0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Table 9: Organic carbon (g kg⁻¹) of rice - growing soil (0-20 cm) at different time interval as influenced by targeted yield- based nutrient application

Treatment	Organic carbon (g kg ⁻¹) at 0-20 cm of rice - growing soil											
	30 DAS			60 DAS			90 DAS			At harvest		
	2013	2014	Mean	2013	2014	Mean	2013	2014	Mean	2013	2014	Mean
RT ₁ - Control	4.90	4.90	4.90	4.80	4.80	4.80	4.80	5.00	4.30	4.50	4.70	4.20
RT ₂ - GRD@ 80:50:30 NPK kg ha ⁻¹	5.30	5.30	5.30	5.50	5.50	5.50	5.50	5.70	5.10	6.10	5.50	5.00
RT ₃ - T.Y.5.0 t ha ⁻¹	5.90	5.80	5.90	6.20	6.20	6.20	5.80	5.60	5.40	6.30	5.60	5.20
RT ₄ - T.Y.6.0 t ha ⁻¹	6.10	6.10	6.10	6.30	6.40	6.40	5.90	6.10	5.90	6.30	5.70	5.40
RT ₅ - T.Y.5.0 t ha ⁻¹ +5 t FYM ha ⁻¹	6.20	6.40	6.30	6.40	6.60	6.50	6.20	5.80	5.90	6.30	5.90	6.10
RT ₆ - T.Y.6.0 t ha ⁻¹ + 5 t FYM ha ⁻¹	6.30	6.50	6.40	6.50	6.60	6.50	6.30	6.40	6.20	6.50	6.40	6.45
S Em ±	0.02	0.01	0.01	0.017	0.008	0.01	0.008	0.020	0.006	0.014	0.011	0.004
CD (p = 0.05)	0.06	0.03	0.04	0.052	0.025	0.03	0.025	0.061	0.020	0.045	0.033	0.014

Table 10: Organic carbon (g kg⁻¹) of wheat - growing soil (0-20 cm) at different time interval as influenced by targeted yield- based nutrient application

Treatment	Organic carbon (g kg ⁻¹) at 0-20 cm of wheat- growing soil											
	30 DAS			60 DAS			90 DAS			At harvest		
	2013-14	2014-15	Mean	2013-14	2014-15	Mean	2013-14	2014-15	Mean	2013-14	2014-15	Mean
WT ₁ - Control	4.33	4.23	4.28	4.35	4.45	4.40	4.73	4.75	4.74	4.68	4.75	4.71
WT ₂ - GRD@ 120:80:60 NPK kg ha ⁻¹	5.20	5.13	5.16	4.80	4.85	4.83	5.25	5.40	5.33	5.28	5.35	5.31
WT ₃ - T.Y.4.5 t ha ⁻¹	5.33	5.43	5.38	4.95	5.13	5.04	5.38	5.60	5.49	5.35	5.53	5.44
WT ₄ - T.Y.6.0 t ha ⁻¹	5.50	5.58	5.54	5.05	5.33	5.19	5.63	5.70	5.66	5.53	5.63	5.58
WT ₅ - T.Y.4.5 t ha ⁻¹ + 5 t FYM ha ⁻¹	5.53	5.63	5.58	5.25	5.50	5.38	5.65	5.73	5.69	5.65	5.75	5.70
WT ₆ - T.Y.6.0 t ha ⁻¹ + 5 t FYM ha ⁻¹	5.55	5.70	5.63	5.35	5.50	5.43	5.90	5.98	5.94	5.85	5.98	5.91
S Em ±	0.009	0.006	0.006	0.006	0.006	0.004	0.006	0.006	0.003	0.010	0.019	0.010
CD (p = 0.05)	0.030	0.018	0.018	0.018	0.018	0.012	0.018	0.018	0.012	0.032	0.058	0.031

Table 11: Effect of soil test based fertilizer application on growth and yield attributes of rice crop (2013-2014)

Treatments	Plant height (cm)		Tillers hill ⁻¹		Panicles hill ⁻¹		Panicle length (cm)	
	2013	2014	2013	2014	2013	2014	2013	2014
RT ₁ - Control	44.85	43.06	4.88	4.35	4.15	4.15	17.45	17.34
RT ₂ - GRD@ 80:50:30 NPK kg ha ⁻¹	63.95	59.91	7.75	7.61	7.47	7.47	20.28	20.79
RT ₃ - T.Y.5.0 t ha ⁻¹	72.80	69.06	8.50	8.43	8.31	8.31	20.95	21.07
RT ₄ - T.Y.6.0 t ha ⁻¹	73.70	70.93	8.98	8.63	8.55	8.55	22.47	23.02
RT ₅ - T.Y.5.0 t ha ⁻¹ +5 t FYM ha ⁻¹	77.70	70.92	9.20	8.97	8.87	8.87	22.02	23.19
RT ₆ - T.Y.6.0 t ha ⁻¹ + 5 t FYM ha ⁻¹	78.50	73.92	9.78	9.29	9.23	9.23	22.40	23.25
S Em ±	2.93	1.62	0.22	0.29	0.32	0.32	0.48	0.36
CD(p=0.05)	8.82	4.90	0.68	0.89	0.97	0.97	1.44	1.08

Table 12: Effect of soil test based fertilizer application on yield and protein content of rice

Treatment	Grain (t ha ⁻¹)		Straw (t ha ⁻¹)		Protein content (%)	
	2013	2014	2013	2014	2013	2014
RT ₁ - Control	3.11	2.21	4.32	3.47	5.80	5.97
RT ₂ - GRD@ 80:50:30 NPK kg ha ⁻¹	4.16	4.26	5.95	6.38	6.63	6.37
RT ₃ - T.Y.5.0 t ha ⁻¹	4.57	4.78	6.53	7.25	6.74	6.61
RT ₄ - T.Y.6.0 t ha ⁻¹	5.94	5.22	8.50	7.97	7.29	7.00
RT ₅ - T.Y.5.0 t ha ⁻¹ +5 t FYM ha ⁻¹	4.79	5.57	6.84	8.53	7.73	7.16
RT ₆ - T.Y.6.0 t ha ⁻¹ + 5 t FYM ha ⁻¹	6.62	5.63	9.47	8.68	8.12	7.70
S Em ±	0.23	0.17	0.33	0.28	0.49	0.185
CD(p=0.05)	0.69	0.52	0.99	0.85	1.46	0.556

References

1. Anonymous. Rice in India hand book of statistics, Director of Rice Development, Government of India, Hyderabad (A.P.), 2006.
2. Aulakh MS. Integrated nutrient management for sustainable crop production, improving crop quality and soil health, and minimizing environmental pollution. : Management and protection of receiving environments, 2010, 79-82.
3. Aulakh MS, Benbi DK. Enhancing fertilizer use efficiency. In: Proceedings of FAI Annual Seminar 2008, 4-6 December, 2008. The Fertilizer Association of India, New Delhi, India. 2008; SII-4:1-23.
4. Dwivedi BS, Pandey AK, Tiwari RK, Jha AK. Performance of integrated nutrient management on yield of paddy. National seminar on Innovation extension approaches for enhancing rural household income. J.N.K.V.V, Jabalpur (M.P.), 2011.
5. Ghosh PK, Bandopadhyay KK, Misra AK, Rae AS. 2004. Balanced fertilization for maintaining soil health and sustainable agriculture. Fert. News. 2011; 49(4):13-35.
6. Gomez KA, Gomez AA. Statistical procedure for agricultural research IInd edition. John Wiley and Sons, New York. Inc. 1984, 680.
7. Hegde DM, Babu SNS. Balanced fertilization for nutritional quality in oilseeds. Fert. News. 2004; 49(4):52-93.
8. Jackson ML. Soil Chemical Analysis. Prentice Hall of India. Pvt. Ltd. New Delhi, 1973.
9. Mishra SK, Tripathi BN, Dixit RN, Singh Ashwini. Production potential of rice-wheat crops, nutrient uptake and soil properties as influenced by fertilizer nutrients and farmyard manure under five years rice-wheat cropping system. Annals of Plant and Soil Research. 2015; 17:193-197.
10. Mishra SK, Tripathi BN, Dixit RN, Singh Ashwini. Production potential of rice-wheat crops, nutrient uptake and soil properties as influenced by fertilizer nutrients and farmyard manure under five years rice-wheat cropping system. Annals of Plant and Soil Research. 2015; 17:193-197.
11. Pandya N, Chouhan GS, Nepalia V. Effect of varieties crop geometric and weed management on nutrient uptake by soybean [*Glycine max* (L.) Merrill] and associated weeds. Indian J of Agronomy. 2005; 50(3):218-220.
12. Pham Sy Tan, Tuyen TQ, Huan TTN, Khuong TQ, Diep LN, Dung HT *et al.* Soil test based fertilizer application on growth and yield of rice: a case study in Omon district, Cantho province. OMON RICE, 2000; 8:64-74.
13. Prasad R, Kumar D, Sharma SN, Gautam RC, Dwivedi MK. Current status and strategies for balanced fertilization. Fert. News. 2004; 49(12):73-80.
14. Ramamoorthy B, Narsimham RL, Dinesh RS. Fertilizer application for specific yield targets of Sonara-64. Indian Farm. 1967; 17:43-45.
15. Roy HK, Kumar A, Sarkar AK, Prasad R, Dubey SC, Kumar A. Yield, nutrient uptake, pest and disease incidence in upland rice as influence by N, K and FYM application in acid sedentary soils. Journal Pot. Research, 1997; 13:131-136
16. Sahu LR, Pandey N, Chandu Lal, Sandeep N, Tiwari BS. Effect of Reference of different agrochemicals in conjunction with different source of nutrient supply on growth characters, yield enhancement and soil fertility. Annals of Plant and Soil Research. 2015; 17:87-90.
17. Subba Rao A, Reddy KS. Implications of soil fertility to meet future demand: Indian scenario. In Proceedings of the IPI-OUAT-IPNI International Symposium on Potassium Role and Benefits in Improving Nutrient Management for Food Production, Quality and Reduced Environmental Damages, (Eds. MS Brar and SS Mukhopadhyay), 5-7 November 2009. IPI, Horgen, Switzerland and IPNI, Norcross, USA. 2009; 1:109-135.
18. Tayefe MA, Gerayzade, Nasrollah A. Effect of nitrogen fertilizer on rice quality. Eco. Env. and Cons. 2010; 16:431-436.
19. Chaubey AK, Parganiha OP, Paraye PM. Effect of STCR technology for targeted yield in rice, 2015.