



E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2018; 7(6): 710-712
Received: 01-09-2018
Accepted: 02-10-2018

Abhiram Dash
Agricultural Statistics, Odisha
University of Agriculture and
Technology, Odisha, India

RK Patnaik
Plant Breeding and Genetics,
Odisha University of Agriculture
and Technology, Odisha, India

SK Dwivedi
Agronomy, Odisha University of
Agriculture and Technology,
Odisha, India

BS Bishoyi
Agronomy, Odisha University of
Agriculture and Technology,
Odisha, India

P Shatpathy
Plant Physiology, Odisha
University of Agriculture and
Technology, Odisha, India

Analysis of rainfall distribution in Bargarh district of Odisha

Abhiram Dash, RK Patnaik, SK Dwivedi, BS Bishoyi and P Shatpathy

Abstract

The variation in the distribution of rainfall over time results in drought, abnormal and normal rainfall in different years. The statistical study of rainfall probability is of utmost important for proper planning of the crops of a region due to the random nature of rainfall pattern. The probability distribution of drought, normal and abnormal rainfall at Bargarh district in western Odisha is studied monthly and seasonally from rainfall data for period of 24 years (1993-94 to 2016-17). The month having monthly rainfall less than half of mean monthly rainfall is regarded as drought month, more than twice the mean monthly rainfall is regarded as abnormal month and in between half and twice of the mean monthly rainfall is regarded as normal month. The probability of normal rainfall remains high during kharif season (June – September) with very low probability occurrences of drought months. The probability of drought months remains high in rabi season, whereas, in summer season the probability of abnormal months remains high. The C.V. of rainfall is less in months having more rainfall and increases in dry months. Thus there is high degree of uncertainty of rainfall in dry months, whereas, the wet months have consistency in their rainfall.

Keywords: Probability, drought months, normal months and abnormal months

Introduction

The success of agriculture of a region depends greatly on the quantity and distribution of rainfall in the region. The random nature of rainfall distribution alongwith its time dependent nature creates a need for its statistical analysis on basis of its probability. Many workers (Sharma *et al*, 1999, Hooda and Thakur, 1998) [4, 2] have studied the occurrence of drought, normal and abnormal rainfall by using probability approach. Drought and floods being natural disasters have direct impact on socioeconomic condition of farmers and also on crop production. Thus it necessitates the study of occurrence of drought, normal and abnormal rainfall of a region. Bargarh district, situated in western part of Odisha has a tropical climate. Rice is the important crop grown in this district whose production is greatly affected by rainfall pattern. In fact, Bargarh district is regarded as the 'Rice Bowl of Odisha'. The study made an effort to examine the probability distribution of drought, normal and abnormal months and seasons at Bargarh district. The seasons are classified as Kharif (June – September), Rabi (October- January) and Summer (February – May).

Material and methods

Monthly rainfall data from 1993-94 to 2016-17 has been collected from IMD, Bhubaneswar. The data are arranged monthly and according to seasons. Each month and season has 24 rainfall events corresponding to 24 years. Monthly and seasonal rainfall events were then classified as drought, normal and abnormal.

Determination of drought, normal and abnormal months

Let $T_1 = \frac{\bar{V}}{2}$; $T_2 = 2\bar{Y}$, where \bar{Y} is the mean monthly rainfall;

If monthly rainfall is less than T_1 , then it is defined as drought month.

If monthly rainfall lies between T_1 and T_2 , then it is defined as normal month.

If monthly recovery rainfall is more than T_2 , then it is defined as abnormal month. (Dash and Shatpathy, 2018) [1].

Percentage of drought, normal and abnormal months is given by

Percentage of drought, normal and abnormal months is given by: (No. of drought, normal and abnormal months / total no. of months in the entire period of study) X 100

Correspondence
Abhiram Dash
Agricultural Statistics, Odisha
University of Agriculture and
Technology, Odisha, India

Probability of occurrence of drought, normal and abnormal months is given by

If a_i, b_i and c_i ($i = 1, 2, \dots, 12$) are the number of drought, normal and abnormal months, out of total number of drought (A), normal(B) and abnormal(C) months during the entire period of 24 years (1993-94 to 2016-17), then the probability of occurrence of drought, normal and abnormal months are given by: $P(A) = a_i/A$; $P(B) = b_i/B$ and $P(C) = c_i/C$; $i= 1, 2, 3, \dots, 12$.

Calculation of probability of occurrence of drought, normal and abnormal seasons is given by

If a_i^*, b_i^* and c_i^* ($i = 1, 2, 3$) are the total number of drought, normal and abnormal months in the i th season, the estimated values of probability of occurrence of drought, normal and abnormal months falling in a given season are given by:

$P(A)^* = a_i^*/A$; $P(B)^* = b_i^*/B$ and $P(C)^* = c_i^*/C$; $i= 1, 2, 3$

The variability in rainfall is studied with the help of Coefficient of variation (Sarkar, *et al.* 2013)^[3].

C.V. of monthly rainfall = $\frac{\text{Standard Deviation of monthly rainfall}}{\text{Mean monthly rainfall}} \times 100$

C.V. of seasonal rainfall = $\frac{\text{Standard Deviation of seasonal rainfall}}{\text{Mean seasonal rainfall}} \times 100$

Correlation coefficient between mean monthly rainfall (X) and C.V. of monthly rainfall (Z) denoted by $r(X,Z)$ is given by $r(X,Z) = \text{Cov}(X, Z)/(\sigma_x \sigma_z)$

Standard error of the Correlation Coefficient is given by $SE(r)$

= $\sqrt{\frac{1-r^2}{n-2}}$, where 'n' is the no. of observations. Here $n = 12$.

Results and discussion

The study of Table 1 which shows the monthly rainfall statistics of Bargarh district reveals that the month of July has the highest mean monthly rainfall and the month of February has the lowest monthly rainfall. This shows that July is the wettest month and February is the driest month over the period of 24 years (1993-94 to 2016-17). Also from table 1 it is found that the wettest month i.e. July has the lowest C.V. and the driest month i.e. February has the highest C.V. Thus an inverse relationship is found to exist between mean and C.V. of monthly rainfall. The correlation coefficient between mean monthly rainfall and C.V. of monthly rainfall is highly negative and significant at 1% level of significance ($r = -0.86^{**}$ with standard error of 0.16). The frequency of normal months is more in wettest months and the frequency of drought months is more in drier months. The probability of normal months is more in wet months and the probability of drought months is more in drier months. The month of December has the highest probability of drought month and the months of July and August has both highest probability of normal month. The probability of abnormal month is highest for the month of April.

The study of seasonal rainfall statistics as revealed from Table 3 shows that the C.V. of seasonal rainfall is lowest for Kharif season and highest for rabi season. Thus it is found that an inverse relationship between mean and C.V. of seasonal rainfall also exist in case of seasons. The probability of normal months is more in kharif season and the probability of drought months is more in rabi season.

Table 1: Monthly Rainfall Statistics of Bargarh District of Odisha


Month	Mean monthly rainfall (in mm)	Mean monthly rainfall as % of mean annual rainfall	C.V.	A ₁	A ₂	No. of drought/normal/abnormal months		
						a _i	b _i	c _i
January	8.72	0.64	216.22	4.36	17.44	18	2	4
February	5.28	0.39	145.61	2.64	10.56	16	6	2
March	7.12	0.52	185.98	3.56	14.24	14	2	8
April	12.91	0.95	155.75	6.46	25.82	16	0	8
May	20.82	1.53	147.61	10.41	41.64	14	6	4
June	184.03	13.55	60.63	92.02	368.06	2	20	2
July	400.92	29.51	38.66	200.46	801.84	0	24	0
August	394.45	29.03	57.78	197.23	788.9	2	20	2
September	227.56	16.75	60.13	113.78	455.12	4	18	2
October	77.95	5.74	125.26	38.98	155.9	10	10	4
November	12.17	0.9	156.03	6.09	24.34	16	2	6
December	7.25	0.53	239.34	3.63	14.5	20	0	4
					Total	132	110	46

Table 2: Probability of drought, normal and abnormal months in a year for Bargarh District of Odisha

Month	Drought month		Normal month		Abnormal month	
	P(A)	Percentage of months having a given month as drought month	P(B)	Percentage of months having a given month as normal month	P(C)	Percentage of months having a given month as abnormal month
January	0.15	75	0.03	8.33	0.13	16.67
February	0.15	66.67	0.04	25	0.06	8.33
March	0.11	58.33	0.04	8.33	0.19	33.33
April	0.07	66.67	0.05	0	0.25	33.33
May	0.05	58.33	0.12	25	0	16.67
June	0.05	8.33	0.11	83.33	0.06	8.33
July	0.02	0	0.15	100	0	0
August	0.02	8.33	0.15	83.33	0	8.33
September	0.02	16.67	0.14	75	0.06	8.33
October	0.03	41.67	0.12	41.67	0.06	16.67
November	0.15	66.67	0.05	8.33	0.06	25
December	0.18	83.33	0	0	0.13	16.67

Table 3: Seasonal Rainfall Statistics and Probability of Drought, Normal and Abnormal Months in a Year for Bargarh District of Odisha

Season	Mean rainfall (in mm)	Mean monthly rainfall as percentage of mean annual rainfall	C.V.	Probability		
				P(A)*	P(B)*	P(C)*
Kharif (June-September)	1210.07	88.11	22.27	0.11	0.51	0.38
Rabi (October-January)	109.01	8.02	89.75	0.55	0.19	0.26
Summer (February-May)	50.14	3.87	99.39	0.13	0.38	0.49

**Fig 1:** Mean and C.V. of Monthly Rainfall in Bargarh District of Odisha**Conclusion**

It is found that the probability of normal months is highest for kharif season, which has highest mean rainfall of 1210.07 mm. Also the C.V. of rainfall in kharif being much lower (22.27 per cent) indicates more consistency in kharif rainfall. The wet months having low C.V. shows consistency in their rainfall pattern whereas the dry months show more variation in their rainfall pattern as the C.V. of rainfall for dry months is very high (99.39). This shows an inverse relationship between mean and C.V. of rainfall. (Fig. 1). The amount of rainfall in kharif season contributing 88.95 per cent to the mean annual rainfall is sufficient for paddy crop. So, paddy can be taken as the main crop in kharif season in Bargarh district. The probability of drought months is highest for the dry months of rabi season, whereas, the wet months i.e., July, August has highest probability of normal months.

References

1. Dash A, Shatpathy P. Probability Analysis of Rainfall Distribution in Sambalpur District of Odisha. *Journal of Pharmacognosy and Phytochemistry*. 2018; 7(5):3169-317.
2. Hooda BK, Thakur BC. Probability Analysis of Rainfall at Naini, Himachal Pradesh. *Indian Journal of Soil Conservation*. 1998; 26(2):153-155.
3. Sarkar S, Kundu DK, Mahapatra BS. Rainfall Probability Analysis of Western Odisha Plateau Region for Sisal (*Agave sisalana perrine eEugelm*) based cropping system. *Journal of Agricultural Physics*. 2013; 13(1):62-70

4. Sharma HC, Chauhan HS, Ram Seva. Probability Analysis of Rainfall for Crop Planning. *Journal of Agri. Engg.* 1999; 17(3):38-44.