



E-ISSN: 2278-4136
P-ISSN: 2349-8234
JPP 2019; 8(3): 881-884
Received: 07-03-2019
Accepted: 09-04-2019

D Leninraja

Assistant Professor (SS&AC) -
Agricultural College & Research
Institute, Tamil Nadu
Agricultural University,
Killikulam, Tuticorin,
Tamil Nadu, India

D Muthumanickam

Professor (SS&AC) – Agriculture
Research Station, Tamil Nadu
Agricultural University,
Bhavanisagar, Tamil Nadu,
India

D Balamurugan

Assistant Professor (SS&AC) -
JKK Munirajah College of
Agricultural Science, Erode,
Tamil Nadu, India

Correspondence**D Leninraja**

Assistant Professor (SS&AC) -
Agricultural College & Research
Institute, Tamil Nadu
Agricultural University,
Killikulam, Tuticorin,
Tamil Nadu, India

Reassessment of soil available DTPA-iron status in Tirunelveli district of Tamil Nadu using GIS and GPS techniques

D Leninraja, D Muthumanickam and D Balamurugan

Abstract

A study was undertaken in Tirunelveli district of Tamil Nadu with a view to assess the DTPA-Fe status of soils at block level. A sum of 1,798 geo-referenced surface soil samples from nineteen blocks of Tirunelveli district representing different soil units as per the soil map prepared on 1:50,000 scales were collected randomly at 0-15 cm depth using Global Positioning System. The soil samples were analysed for available DTPA-Fe. The content of DTPA- Fe differed from 0.05 to 72.31 mg kg⁻¹ soil. Analytical results and the GPS data were used for the preparation of thematic maps showing spatial distribution of micronutrients status block wise in the district. Locations of soil sampling sites of Tirunelveli district were marked on base map on 1: 50,000 scales prepared from State Revenue Maps and digitized using Arc-info GIS. The delineation study thus clearly indicates that, DTPA-Fe was found to be deficient in 49.9 per cent of soils of Tirunelveli district respectively.

Keywords: DTPA-Fe, GPS, GIS, thematic maps

1. Introduction

Soil micronutrients play a vital role in the growth, development and yield of plant besides the information on the nutritional status of an area and thus go a long way in planning judicious fertilizers and soil management practices to develop economically viable alternatives for farming community. The estimation, characterization and comparison of micronutrients of soil are important issues in site-specific crop management, precision farming and sustainable agriculture (Deb, 1997) [1].

In the context of today changing scenario, there is a need to generate the spatial data of micronutrients using frontier technologies like Global Positioning System (GPS) and Geographical Information System (GIS). The GPS has revolutionized positioning concept though it started primarily as a satellite based radio navigation system providing precise, three dimensional position navigation and time information. The GIS provides scientists, planners, managers and decision makers an efficient way of combining and analyzing geo-referenced and descriptive data from different sources (soils, vegetation, geology, land covers and others) for better understanding and management of natural resources (Fernandez *et al.* 1993) [2]. The thematic maps for individual nutrient (Zn, Fe, Cu and Mn) is prepared by using GIS software (Nayak *et al.*, 2006) [6] and multi micronutrient maps are generated by integrating individual maps of Fe, Mn, Zn and Cu in the GIS (Sood *et al.* 2004) [7]. This will also help in monitoring changes in micronutrient status over a period of time. It can be revisited with help of GPS, which is otherwise not possible in the random sampling.

With this background a study was conducted with revolutionary effort to examine soil available micronutrient status and delineate the available micronutrients status scrupulously at block level in the Tirunelveli district of Tamil Nadu.

1.1 Study Area

Tirunelveli district in Tamil Nadu is bounded by Virudhunagar district in the North, Western Ghats in the West, Kanniyakumari district in the South and Thoothukudi district in the East. Tirunelveli district is comprised of 11 taluks, 19 blocks and 628 Revenue villages covering an area of 6, 81,065 ha of land. The Tirunelveli district lies between 8°.08' and 9°.25' of the Northern latitude and 77°.09' and 77°.59' of Eastern longitude. Major portion of the district is covered by plain topography. Red loam is the predominant soil type in the district accounting for 48.21 per cent followed by the black soil of 30.09 per cent.

2. Material and Methods

2.1 Collection of Soil Samples

Totally 1798 geo-referenced surface soil samples covering the entire village in nineteen blocks of Tirunelveli district were collected randomly at 0-15 cm depth by adopting the standard procedures of soil sample collection. The GPS data (Latitude °N and Longitude °E) were collected from each sampling sites distributed over the entire Tirunelveli district by using Garmin

GPS 76CS model (Fig. 1). The collected soil samples were dried, gently bound, sieved (2 mm sieve) and preserved in polythene bags for DTPA extractable micronutrients (Lindsay and Norvell, 1978) [4]. Locations of soil sampling sites of Tirunelveli district were marked on base map on 1: 50,000 scales prepared from State Revenue Maps and digitized using Arc-info GIS (9.2).




Fig 1: Soil sampling points of Tirunelveli district

2.2 Generation of Map

The Tirunelveli district map (1:50,000) was map was vectorised by using Raster to Vector software (R2V), and then exported into Arc-GIS software. Database on soil micronutrient status of the study area was developed using Microsoft Excel package. The database was exported to Arc GIS software and the thematic maps on different available micronutrients status were generated. The thematic maps at block level were generated for showing status of different available micronutrients status based on block mean and nutrient index values.

3. Results and Discussion

In Tirunelveli district, the two major soil groups exists are the red and black soils. Red loam is the predominant soil type accounting for 48.21 per cent followed by the black soil of 30.09 per cent. The other types of soils are lateritic soil, sandy coastal alluvium, red sandy soil and others.

3.1 DTPA- Iron (Table 1)

The content of DTPA- Fe differed from 0.05 to 72.31 mg kg⁻¹ with an average of 7.74 mg kg⁻¹ soil. Considering 4.5 mg kg⁻¹ as critical limit as suggested by Lindsay and Norvell (1978) [4] the soils of this district had sufficient amount of available iron. Among the blocks studied, Kunnathur village of Palayamkottai block showed the highest value of 72.31 mg kg⁻¹ with respect to available Fe followed by Brammadesam of Ambasamudram block as (54.14 mg kg⁻¹). Since the soils of Palayamkottai and Ambasamudram blocks are noncalcareous in nature due to that the DTPA-Fe content was found to be high (Murthy *et al.* 1997) [5] while the soils of Sangupatti village of Kuruvikulam block showed the lowest mean value (0.05 mg kg⁻¹) of available Fe content. The CaCO₃ content has negative influence on the DTPA extractable iron content in that block. Deficiency of iron in the calcareous soils is well documented (Lindsay and Norwell 1978) [4].

Table 1: Range, Mean values and Percent Sample Category of DTPA-Fe status for different blocks of Tirunelveli district

S. No	Block name	Range and Mean Values Fe (mg kg ⁻¹)	Percentage sample category		
			DTPA- Fe		
			L	M	H
1	<u>Alangulam</u>	0.09 - 20.58 (3.31)	64.8	29.6	5.6
2	<u>Ambasamudram</u>	10.81 - 54.14 (34.38)	0.0	0.0	100.0
3	<u>Cheranmahadevi</u>	0.93 - 6.87 (2.51)	78.6	21.4	0.0
4	<u>Kadayanallur</u>	1.41 - 22.91 (5.28)	50.7	29.9	19.4
5	<u>Kadayam</u>	0.86 - 27.28 (8.32)	34.1	31.8	34.1
6	<u>Kalakadu</u>	0.85 - 25.57 (7.41)	46.2	15.4	38.5
7	<u>Keelapavoor</u>	0.32 - 5.82 (1.86)	92.9	7.1	0.0
8	<u>Kuruvikulam</u>	0.05 - 39.33 (6.42)	26.4	24.1	49.4
9	<u>Manur</u>	0.40 - 35.51 (6.85)	52.6	22.4	25.0
10	<u>Melaneelithanallur</u>	0.10 - 22.8 (2.69)	75.0	22.0	3.0
11	<u>Nanguneri</u>	0.95 - 19.23 (5.57)	40.5	34.5	25.0
12	<u>Pappakudi</u>	2.46 - 46.29 (9.56)	11.3	56.5	32.3
13	<u>Palayamkottai</u>	2.74 - 72.31 (23.17)	13.2	29.5	57.4
14	<u>Radhapuram</u>	0.37 - 4.07 (1.52)	97.1	2.9	0.0
15	<u>Sankarankovil</u>	0.09 - 19.58 (2.71)	82.8	7.8	9.5
16	<u>Senkottai</u>	1.53 - 29.69 (9.48)	0.0	0.0	100.0
17	<u>Tenkasi</u>	0.28 - 43.59 (15.57)	24.7	27.3	48.1
18	<u>Valliyoor</u>	0.72 - 18.03 (5.58)	61.3	12.0	26.7
19	<u>Vasudevanallur</u>	0.89 - 35.74 (4.09)	65.2	20.7	14.1
		Overall for District	49.9	21.4	28.6

3.2 Thematic maps

The DTPA extractable iron of different blocks of Tirunelveli district was grouped into three categories based on the critical limits followed for availability of micronutrient in India and Tamil Nadu. The thematic map clearly identifies the blocks that are extremely deficient in micronutrient status which

require utmost attention to sustain the soil productivity. Out of nineteen blocks, six blocks recorded high DTPA-Fe status (Fig.2). The sufficient level of iron status was attributed to the origin of soil, in which 48.21% of the soils in Tirunelveli district are red loamy soil.




Fig 2: DTPA- Fe status of Tirunelveli District

4. Conclusion

Thus, from the above investigation it is very clear that the soils of Tirunelveli district are moderately deficient in available iron. Therefore, to overcome the deficiency of iron which are tending towards increasing deficiency, the need for sustained application of organic manures or through supplementation with inorganic fertilizers is essential. Besides, the edaphic factors such as organic carbon content and free lime status are widely believed to be circumventing the availability of micronutrients, which also needs to be addressed to develop strategies for alleviation of micronutrient deficiencies in Tirunelveli district of Tamil Nadu. Micronutrient status plays a major role in increasing crop yields and soil productivity in general and hence, it is essential to adopt an integrated way of adding organic amendments with micronutrients, which will sustain the soil fertility and crop productivity.

5. References

1. Deb DL. Micronutrient research and crop production in India. *Journal of the Indian Society of Soil Science*. 1997; 45:675-692.
2. Fernandez RN, Rusinkiwicz M, Morais da Silva L, Johannsen CJ. Design and Implementation of soil geographic database for rural planning and management. *Journal of soil and water conservation*. 1993; 48:140-145.
3. Krishnasamy R, Shanmugam M, Poongothai S, Valliappan K, Mani S, Mallika Vanangamudi, Santhi P, Devarajan R. Twenty five years of micronutrients research in soils and crops of Tamil Nadu 1967-1992. *Research Bulletin, Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore- 3*, 1994.
4. Lindsay NL, Norvell WA. Development of DTPA soil test for zinc, iron, manganese and copper. *Soil Science Society of America*. 1978; 42:421-428.
5. Murthy IY, Sastry LN, Datta TG, Narayanasamy G, Rattan RK. Distribution of micronutrient cations in vertisols derived from different parent materials. *Journal of the Indian Society of Soil Science*. 1997; 45:577-580.
6. Nayak AK, Chyinchamalathpure, Anil R, Gururaja Rao, G, Khandelwal MK, Tyagi NK. Spatial variability of DTPA extractable micronutrients in soils of Bara tract of Sardar Sarovar canal command in Gujarat state India. *Journal of the Indian Society of Soil Science*. 2006; 42:137-145.
7. Sood Anil, Setia RK, Bansal RL, Sharma PK, Nayyar V. Spatial variability of micronutrients in soils of Amritsar district using frontier technologies. In: *Proceedings of 7th Punjab Science Congress*. February 7- held at Guru Nanak Dev. University, Amritsar, 2004.
8. Takkar PN, Nayyar VK, Bansal RL, Dwivedi RS, Manna MS. Annual progress report of ICAR co-ordinated micronutrient scheme 1996-97. *Punjab Agricultural University, Ludhiana*, 1997.