



E-ISSN: 2278-4136

P-ISSN: 2349-8234

[www.phytojournal.com](http://www.phytojournal.com)

JPP 2020; 9(2): 1621-1623

Received: 04-01-2020

Accepted: 06-02-2020

**S Muthuramu**Agricultural Research Station,  
Tamil Nadu Agricultural  
University, Paramakudi,  
Tamil Nadu, India**T Ragavan**Agricultural Research Station,  
Tamil Nadu Agricultural  
University, Paramakudi,  
Tamil Nadu, India

## Genotypic correlation and path coefficient analysis for yield traits in rainfed rice (*Oryza sativa* L.)

**S Muthuramu and T Ragavan**

**Abstract**

Genotypic correlation and path coefficient analysis was carried out at Agricultural Research Station, Tamil Nadu Agricultural University, Paramakudi to study the association between nine yield related traits during Rabi 2018-19 in 38 rice genotypes under rainfed rice ecosystem. Number of productive tillers per plant, number of panicles per square metre, straw yield per plot and harvest index expressed positive and significant correlation with grain yield. The trait, number of productive tillers per plant had positive and significant association with number of panicles per square metre, spikelet fertility and harvest index suggesting that selection based on number of productive tillers per plant is highly fruitful in developing high yielding genotypes in rainfed situation, as it will bring simultaneous improvement of these traits. The traits straw yield and harvest index expressed high direct effect on grain yield. The traits number of productive tillers per plant, number of panicles per square metre, straw yield per plot and harvest index should be given more importance for enhancing grain yield under rainfed rice ecosystem.

**Keywords:** *Oryza sativa*, rainfed, genotypic correlation, path coefficient

**Introduction**

Rice (*Oryza sativa* L.) is one of the major cereal food crops for more than 50 per cent of the world population. It is cultivated over 167 million hectares with the production of 780 million tonnes (FAOSTAT, 2017). In India rice is cultivated in an area of 44.50 million hectares with a production of 116.00 million metric tons during 2018-19 (USDA 2019). Grain yield in rice is a quantitatively inherited trait and involves function of several components. Selection of superior genotypes based on yield is difficult due to the integrated structure of plant in which the component characters are interdependent and are governed by a large number of genes.

An idea on the extent of association between traits conferring higher yield will be much helpful to decide upon the traits to be given importance in selection process. A positive association between traits warrants the simultaneous improvement of both the traits while restricting selection to any one of the associated traits. On the other hand, a negative relationship between two traits necessitates equal weightage to be given on both the traits during selection. At genetic level, a positive correlation occurs due to coupling phase of linkage and negative correlation arises due to repulsion phase of linkage of genes controlling two different traits (Nadarajan and Gunasekaran, 2008) [10]

Path analysis has been used to organize the relationship between predictor variables and response variables. The advantage of path analysis is that it permits the partitioning of the correlation coefficient into its components-one component being the path coefficient (or standardized partial regression coefficient) that measures the direct effect of a predictor variable upon its response variables, the second component being the indirect effect(s) of a predictor variable on the response variable through other predictor variables (Dewey and Lu, 1959) [3]. Path coefficient analysis assists plant breeders in identifying traits on which selection pressure should be given for improving yield. With these points in view, the present study was framed to study the relationship between yield related traits under rainfed rice ecosystem.

**Materials and Methods**

The experimental material comprised with thirty eight advanced rice cultures which were evaluated in a randomized block design with three replications at Agricultural Research Station, Tamil Nadu Agricultural University, Paramakudi during Rabi 2018-19. The experimental site is located at 9° 21' N latitude, 78° 22' E longitudes and an altitude of 242 m above mean sea level with average annual rainfall of 840 mm. This site has clay loam soil texture with pH of 8.0. Each genotype was raised in 5x2 m plot keeping 15 x 10 cm spacing. The recommended agronomic practices followed to raise good crop stand.

**Corresponding Author:****S Muthuramu**Agricultural Research Station,  
Tamil Nadu Agricultural  
University, Paramakudi,  
Tamil Nadu, India

The data were recorded on ten randomly selected plants from each replication for various quantitative traits studied were *viz.*, days to 50% flowering, plant height (cm), number of productive tillers per plant, number of panicles per square metre plot area, panicle length (cm), number of filled grains per panicle, spikelet fertility (%), grain yield per plot (kg), straw yield per plot (kg) and harvest index (%).

The genotypic correlation coefficients between yield and yield components as well as among the yield components were worked out. From the analysis of variance and covariance tables, the corresponding genotypic variances and covariances were calculated by using the mean square values and mean sum of products as suggested by Al-Jibouri *et al.* (1958). The relative influence of eight components on yield by themselves (direct effects) and through other traits (indirect effects) was evaluated by the method of path coefficient analysis as suggested by Dewey and Lu (1959) [3]. The simple correlation coefficients already estimated at genotypic level were utilized for this purpose. By keeping yield as dependent variable and other eight traits as independent variables, simultaneous equations which express the basic relationship between path coefficients were solved to estimate the direct and indirect effects. The direct and indirect effects were classified based on the scale given by Lenka and Misra (1973) [8].

### Results and Discussion

Selection based on the detailed knowledge of magnitude and direction of association between yield and its attributes is very important in identifying the key characters, which can be exploited for crop improvement through suitable breeding programme. Genotypic correlations between yield and yield components *viz.*, days to 50% flowering, plant height, number of productive tillers per plant, number of panicles per square metre, panicle length, number of filled grains per panicle, spikelet fertility (%), straw yield per plot and harvest index (%) were computed for rice genotypes. The results are presented in Table 1.

In the present study, grain yield exhibited positive and significant association with number of productive tillers per plant, number of panicles per square metre, straw yield per plot and harvest index (Table 1). This was in conformity with the findings of Babu *et al.* (2012) [2], Vanisree *et al.* (2013) [13], Islam *et al.* (2015) [6] and Shinde *et al.* (2015) [11], Muthuramu and Sakthivel (2016) [9] and Lalitha *et al.* (2019) [7]. Knowledge on inter relationship between yield traits may facilitate breeder to decide upon the intensity and direction of selection pressure to be given on related traits for the simultaneous improvement of these traits. In the present investigation, number of productive tillers per plant had positive and significant association with number of panicles per square metre, spikelet fertility and harvest index. The trait, plant height showed positive and significant association with panicle length and straw yield per plot. Similarly panicle

length had positive and significant relationship with number of filled grains per panicle and straw yield per plot. The trait, days to 50% flowering showed positive and significant association with number of filled grains per panicle. Likewise positive and significant relationship was observed between number of panicles per square metre and straw yield per plot. Similar results were reported by Babu *et al.* (2012) [2], Islam *et al.* (2015) [6] and Lalitha *et al.* (2019) [7].

Plant height had negative and significant correlation with number of productive tillers per plant, spikelet fertility and harvest index. Number of productive tillers per plant showed negative and significant association with panicle length and number of filled grains per panicle. This was in conformity with the findings of Babu *et al.* (2012) [2] and Muthuramu and Sakthivel (2016) [9].

In the light of above discussion, it may be suggested that, number of productive tillers per plant, number of panicles per square metre, straw yield per plot and harvest index were to be given importance while selection as they expressed positive and significant correlation with grain yield. The trait, number of productive tillers per plant had positively correlated with number of panicles per square metre, spikelet fertility and harvest index suggesting that selection based on number of productive tillers per plant is highly fruitful in developing high yielding genotypes under rainfed condition, as it will bring simultaneous improvement of these traits. A productive tiller per plant is one of the important characters to be considered in the selection for improvement of drought tolerant rice genotypes (Hossain *et al.*, 2018) [5]. Hence result obtained from this study is similar to those mentioned above.

Rapid improvement in yield is expected to result if selection is practiced for component characters. Rate of improvement is expected to be rapid if differential emphasis is laid on the component characters during selection. The basis of differential emphasis could be the degree of influence of component characters on the character of interest. Path analysis gives an idea about how a trait influences grain yield directly and indirectly *via* other traits. This is very important in giving due weightage to major yield contributing traits while selection.

In the present study, straw yield and harvest index expressed high direct effect on grain yield (Table 2). This was in conformity with the findings of Babu *et al.* (2012) [2], Vanisree *et al.* (2013) [13], Islam *et al.* (2015) [6], Shinde *et al.* (2015) [11] and Muthuramu and Sakthivel (2016) [9]. Genotypic residual effect (0.15) indicates that traits under study contribute 85% to the variability in grain yield. It indicates that few other traits which have not been studied here, need to be included to account fully for the variation in grain yield. As a conclusion, number of productive tillers per plant, number of panicles per square metre, straw yield per plot and harvest index should be given more importance for enhancing grain yield under rainfed rice cultivation system.

**Table 1:** Genotypic correlation coefficients for yield related traits and grain yield.

Characters	Days to 50% flowering	Plant Height	Productive tillers per plant	Panicles per square metre	Panicle length	Filled grains per panicle	Spikelet Fertility	Straw yield per plot	Harvest index	Grain yield per plot
Days to 50% flowering	1.000	-0.239	0.087	-0.184	-0.097	0.474**	0.258	-0.245	0.315	0.058
Plant Height		1.000	-0.306*	-0.173	0.772**	0.149	-0.353*	0.449**	-0.326*	0.119
Productive tillers per plant			1.000	0.309*	-0.511*	-0.306*	0.397**	0.025	0.504**	0.648**
Panicles per square metre				1.000	-0.205	-0.285*	0.239	0.495**	-0.131	0.357*
Panicle length					1.000	0.344*	-0.237	0.322*	-0.269*	-0.046
Filled grains per panicle						1.000	0.184	-0.145	0.049	-0.168
Spikelet Fertility							1.000	0.236	0.037	0.263
Straw yield per plot								1.000	-0.605	0.373**
Harvest index									1.000	0.493**
Grain yield per plot										1.000

\*Significance at 5% level; \*\* Significance at 1 % level

**Table 2:** Direct and indirect effects of yield related traits on grain yield

Characters	Days to 50% flowering	Plant Height	Productive tillers per plant	Panicles per square metre	Panicle length	Filled grains per panicle	Spikelet Fertility	Straw yield per plot	Harvest index	Grain yield per plot
Days to 50% flowering	-0.015	0.004	-0.001	0.003	0.002	-0.007	-0.004	0.004	-0.005	0.058
Plant Height	-0.040	0.166	-0.051	-0.029	0.128	0.025	-0.058	0.074	-0.054	0.119
Productive tillers per plant	-0.001	0.002	-0.006	-0.002	0.003	0.002	-0.002	-0.001	-0.003	0.648**
Panicles per square metre	0.009	0.009	-0.016	-0.051	0.011	0.015	-0.012	-0.025	0.007	0.357*
Panicle length	0.022	-0.170	0.113	0.045	-0.221	-0.076	0.052	-0.071	0.060	-0.046
Filled grains per panicle	-0.011	-0.004	0.007	0.007	-0.008	-0.024	-0.004	0.004	-0.001	-0.168
Spikelet Fertility	-0.001	0.002	-0.002	-0.001	0.001	-0.001	-0.005	-0.001	-0.001	0.263
Straw yield per plot	-0.265	0.485	0.027	0.535	0.348	-0.156	0.255	0.898	-0.654	0.373**
Harvest index	0.360	-0.373	0.576	-0.150	-0.309	0.056	0.043	-0.692	0.999	0.493**

\*\* Significance at 1 % level \*\* Significance at 1 % level Residual effect = 0.15 Diagonal values (in bold) denote the direct effects

## References

- Al-Jibouri HR, Miller PA, Robinson HF. Genotypic and environmental variances and covariances in aneuploid cotton cross of inter specific origin. *Agronomy Journal* 1958; 50:633-636.
- Babu VR, Shreya K, Dangi KS, Usharani G, Shankar AS. Correlation and Path Analysis studies in Popular Rice Hybrids of India. *Int. J. Sci. and Res. Publ.* 2012; 2(3):1-5.
- Dewey DR, Lu KH. A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agronomy Journal* 1959; 51:515-518.
- FAOSTAT.2017. Available at [http://faostat3.fao.org/faostatgateway/go/to/download / Q/QC/E](http://faostat3.fao.org/faostatgateway/go/to/download/Q/QC/E). FAO, Rome, Italy.
- Hossain S, Salim M, Golam AM, Noman S. Variability, correlation and path analysis in drought tolerant rice (*Oryza sativa* L.). *J Biosci. Agric. Res.* 2018; 18(2):1521-1530.
- Islam MA, Raffi SA, Hossain MA, Hasan AK. Character association and path coefficient analysis of grain yield and yield related traits in some promising early to medium duration rice advanced lines. *Int. J Expt. Agric.* 2015; 5(1):8-12.
- Lalitha R, Mothilal A, Arunachalam P, Senthil N, Hemalatha G. Genetic variability, correlation and path analysis of grain yield, grain quality and its associated traits in EMS derived M4 generation mutants of rice (*Oryza sativa* L.). *Electronic Journal of Plant Breeding*, 2019; 10(3):1140-1147.
- Lenka D, Misra B. Path co-efficient analysis of yield in rice varieties. *Ind. J Agric. Sci.* 1973; 43:376-379.
- Muthuramu S, Sakthivel S. Correlation and Path Analysis for Yield Traits in Upland Rice (*Oryza sativa*). *Research Journal of Agricultural Sciences.* 2016; 7(4, 5):763-765.
- Nadarajan N, Gunasekaran M. *Quantitative Genetics and Biometrical Techniques in Plant Breeding.* Kalyani Publishers, Ludhiana, 2008, 207.
- Shinde SR, Kumar K, Pawar RM. Genetic studies in upland rice (*Oryza sativa* L.). *Int. J Plant Sci.* 2015; 10(1):33-37.
- United States Department of Agriculture (USDA). 2019. USDA revises India rice and corn production estimates higher. *Agriculture commodities*, [www.indoasiancommodities.com](http://www.indoasiancommodities.com)
- Vanisree S, Swapna K, Raju D, Raju S, Sreedhar M. Genetic variability and selection criteria in rice. *J biological and scientific opinion.* 2013; 1(4):341-346