



E-ISSN: 2278-4136

P-ISSN: 2349-8234

www.phytojournal.com

JPP 2020; 9(4): 1567-1571

Received: 08-05-2020

Accepted: 12-06-2020

Aditi A Deshmukh

Department of Horticulture,
Vegetable Science, Dr.
Panjabrao Deshmukh Krishi
Vidyapeeth, Akola,
Maharashtra, India

Dr. PK Nagre

Department of Horticulture,
Vegetable Science, Dr.
Panjabrao Deshmukh Krishi
Vidyapeeth, Akola,
Maharashtra, India

Dr. AP Wagh

Department of Horticulture,
Vegetable Science, Dr.
Panjabrao Deshmukh Krishi
Vidyapeeth, Akola,
Maharashtra, India

Corresponding Author:**Aditi A Deshmukh**

Department of Horticulture,
Vegetable Science, Dr.
Panjabrao Deshmukh Krishi
Vidyapeeth, Akola,
Maharashtra, India

Effect of nitrogen and phosphorus levels on yield and quality of fenugreek (*Trigonella-foenum-graecum*)

Aditi A Deshmukh, Dr. PK Nagre and Dr. AP Wagh

Abstract

An experiment was conducted during the *rabi* season of 2014-15, to study the "Effect of nitrogen and phosphorus on growth and seed yield of fenugreek" at the Main garden, University Department of Horticulture, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola. The experiment was laid out in Factorial Randomized Block Design with three replications. Twelve treatment combinations were formed with a view to integrate four nitrogen doses and three phosphorus doses. The allocation of treatments was made by random method. On the basis of results obtained in the present investigation, the yield parameters in respect to number of pods per plant, seeds per pod, seed yield per plant, seed yield per plot, seed yield per hectare and test weight were observed significantly maximum in the treatment T₉ (N₃P₃) i.e. 80kg N+60kg P. Similarly, better seed quality parameters *viz.*, germination percentage was recorded significantly with treatment T₁₂(N₄P₃) i.e. 100 kg N + 60 kg P and protein content was recorded significantly superior with treatment T₉ (N₃P₃) i.e. 80 kg N + 60 kg P

Keywords: Fenugreek, yield parameters, test weight, germination percent and protein content nitrogen, phosphorus and yield

Introduction

Fenugreek (*Trigonella foenum-graecum* Linn.) is an important seed spice specially known as methi. It is an annual herb of leguminosae family. Fenugreek is the third largest seed spice in India after coriander and cumin (NBH2011-12). Fenugreek is an important condiment crop grown in Southern India during the kharif and rabi seasons. In India, fenugreek seed production is being under taken on 115.6 thousand hectares area with an annual production of 136.6 thousand tonnes and having productivity of 1.2 tonnes ha⁻¹. The Maharashtra occupies 1122 hectares of area under fenugreek cultivation with an annual production of 4519 tonnes and having productivity of 4.02 tonnes ha⁻¹. In Vidarbha region it is being cultivated on an area of 278 ha with annual production of 1134 tonnes and average productivity of 5.77 tonnes ha⁻¹ (Anon., 2014) [1]. Fenugreek is a rich source of wide variety of components. Fenugreek seeds substantially contain 'Diosgenin' which is used as a starting material in the synthesis of sex hormones. The content of diosgenin in fenugreek seeds varies from 0.40-1.26%. Minerals like Ca (360.0 mg), P (51.0 mg), Thiamine (0.05 mg), Riboflavin (0.15 mg), Na (76.0 mg), K (31.0 mg), Cu (0.26 mg) and Mg (67.0 mg per 100 g fresh weight) are occurred abundantly in fenugreek. It is also a rich source of Vitamin A and C (Das, 1992) [3]. Medicinal value of fenugreek is well recognized in India since antiquity. Methi seeds and leaves are important particularly against the digestive disorders (Sheoran *et al.* 1999) [16] and useful for diabetic patients. Fenugreek seeds are mainly used as a spice for the preparation of different tasty dishes. It also have a high medicinal and industrial importance. It prevents constipation, removes indigestion, stimulates spleen and liver, and is appetizing and diuretic. Seeds are of industrial importance as used for dye and for extraction of alkaloids or steroids. It is also used as a fodder. Indian women use to consume the seeds of fenugreek for its power to promote lactation.

Generally, seed production of methi is taken after 2-3 cuttings, but seed yield obtained without cuttings are better than the seed yield obtained from 2-3 cuttings. It is, therefore, recommended to take seed production of methi without any cuttings (Gill and Singh, 1988). In Maharashtra, though methi is cultivated as an important leafy vegetable and is also grown as a spice, but less attention is being paid on its commercial seed production. Scientific seed production resembles the importance of quality seed to be used for raising of crop in order to get the higher production of good quality. The less availability of quality seeds has been one of the drawback in getting an increased agricultural production in vegetable crops. The availability of pure and good quality seeds has a significance in vegetables seed production; where, more uniformity of

The Colour, size and purity is required. Although, the seed production in vegetables are highly profitable, yet the growers are afraid to take up this venture because of lack of technical knowledge in production technology. Fenugreek seed production is a highly specialized job and it requires intimate knowledge of crop production particularly, the floral biology, mode of pollination, isolation, climatic requirement and nutritional requirements.

Materials and Methods

The field experiment was conducted during the *rabi* season of 2014-15, to study the "Effect of nitrogen and phosphorus on growth and seed yield of fenugreek" at the Main garden, University Department of Horticulture, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola. Akola is situated in the sub-tropical zone at the latitude of 22°42'N and longitude of 77°02'E. The altitude of the place is 307.4 m from mean sea level. The experiment was laid out in Factorial Randomized Block Design with 12 treatments replicated for three times. The layout consisted of 36 plots. The treatments consisted of two factors, nitrogen doses and phosphorus doses. In first factor there are four different nitrogen doses *viz.* N₁ (40kg ha⁻¹), N₂ (60kg ha⁻¹), N₃ (80kg ha⁻¹) and N₄ (100kg ha⁻¹). In second factor there different phosphorus doses *viz.* P₁ (20kg ha⁻¹), P₂ (40kg ha⁻¹) and P₃ (60kg ha⁻¹). Thus in all twelve treatment combinations were studied to find out the suitable combination of nitrogen and phosphorus doses for maximum production of fenugreek. The sowing was done by flat bed system on 24th of December 2014 with spacing of 30cm between rows and 10 cm between plants. The gross and net plot size were 1.8 m X 1.8 m (3.24 m²) and 1.2 m x 1.7 m (2.04 m²).

The soil type of experimental plot was medium black, clay loam and well drained. It was low in available nitrogen (146 kg ha⁻¹), medium in phosphorus (16 kg ha⁻¹) and high in potassium (290 kg ha⁻¹). It was alkaline in reaction (pH 8.1) having EC (0.45 dSm⁻¹). The fenugreek cv. Pusa Early Bunching recommended for all states of northern India, released from CCSHAU, Hisar. It is a mid-late variety with medium to small, oval shaped seeds. Seeds having a light red colour at the time of maturity, dual purpose variety grown for both spices as well as leaves, spreading in nature and resistant to water lodging. It is a high yielding cv. average yield is 20-25q/ha. The doses of nitrogen fertilizer given through use of urea (46% N) and phosphorus through single super phosphate (16% P₂O₅). The experimental field was irrigated immediately after sowing and subsequent irrigations were given to the plots at an interval of 3-4 days during the period of experimentation. Observations were recorded on different parameters, *viz.*, Plant height(cm), No. of branches, Days required for harvesting, Days required for first flowering, Days required for 50% flowering, No. of pods per plant, No. of seeds per pod, Seed yield per plant(g), Seed yield per plot(kg) and Seed yield per hectare(q). The data were statistically analyzed as per the method prescribed and suggested by Panse and Sukhatme (1985) [10].

Results and Discussion

Number of pods per plant, Seeds per pod and seed yield Effect of nitrogen levels

A perusal of data on (Table 1) revealed that number of pods per plant, seeds per pod and seed yield were significantly influenced by different levels of nitrogen at different growth stages. In case of number of pods per plant the data revealed that maximum number of pods (27.33) per plant were recorded with treatment 80 kg Nha⁻¹ which was found to be superior over all other treatments. However, the minimum

number of pod per plant (13.14) were recorded with 40 kg Nha⁻¹. And in case of number of seeds per pod, the maximum seeds per pod (13.64) were recorded with 80 Kg Nha⁻¹. Whereas, the minimum number of seeds per pod (9.01) were observed with application of 40 kg Nha⁻¹. The nitrogen levels show significant increase in seeds per pod. Up to 80 kg N ha⁻¹. In case Seed yield significantly maximum seed yield per hectare (13.27q) was recorded with 80 kg Nha⁻¹, which was found to be significantly superior over other treatments. Whereas, minimum seed yield per hectare (8.24q) was recorded with 40 kg Nha⁻¹. This was probably due to application of higher nitrogen doses resulted in better growth and good seed yield which increased seed yield per hectare. Increased nitrogen level found to have more number of pods per plant and seeds per pod as yield contributing characters. It might be due to fact that, profused nodulation leading to increased nitrogen fixation, which might in turn had profused effect on photosynthetic activity of plant. It accumulates food materials in seed. The results of this investigation supports the results of Patel *et al.* (2010) [12], Chaudhary *et al.* (2011) [4], and Mehta *et al.* (2012) [7] in fenugreek.

Effect of phosphorus

A perusal of data on (Table 1) also revealed that number of pods per plant, seeds per pod and seed yield were significantly influenced by different levels of phosphorus at different growth stages. The significantly maximum pods per plant (18.47) were recorded with 60 kg Pha⁻¹ which was superior among all other treatments. Whereas, minimum number of pods (17.24) were recorded with treatment 20 kg Pha⁻¹. Significantly maximum seeds per pod (11.65) were recorded with 60kgPha⁻¹. Whereas, least number of seeds per pod (10.61) were recorded with treatment 20 Kg Pha⁻¹. Seed yield per hectare was significantly influenced by different phosphorus levels. Significantly maximum seed yield per hectare (11.96q) was recorded with 60 kg Pha⁻¹. However, minimum seed yield per hectare (10.15q) was recorded with 20 kg Pha⁻¹. The seed yield per plant increased with increase of phosphorus application may be due to improved nutritional condition. The higher uptake of nitrogen and phosphorus during seed formation might have increased number of seeds per plant. There was a significant increase in dry matter accumulation at different growth stage which finally reflected in significant improvement in productivity of crop. Significantly maximum seed yield per plant was recorded with 60 kg P ha⁻¹ and the minimum was recorded with 20 kg P ha⁻¹. Similar results were observed by Sheron *et al.* (1999) [16], Jat *et al.* (2001) [5] and Patel *et al.* (2010) [12].

Interaction effect

Interaction effect due to different nitrogen and phosphorus levels on number of pods per plant, seeds per pod and seed yield were found to be significant. The maximum pods per plant (22.77) was recorded from treatment N₃P₃ i. e. 80 kg N and 60 kg P per hectare. Minimum pods per plant (12.51) was recorded with treatment combination N₁P₁ i.e.40 kg N and 20 kg P per hectare. The maximum seed per pod (14.23) was recorded from treatment N₃P₃ i. e. 80 kg N and 60 kg P per hectare. Interaction effect due to different nitrogen and phosphorus levels on seed yield per hectare was found to be significant. The significantly maximum seed yield per hectare (14.35q) was received from the treatment N₃P₃ i. e. 80 kg N and 60 kg P per hectare. Whereas, minimum yield per hectare (6.96q) was recorded from treatment of N₁P₁. The results of this investigation supports the results of Mavai *et al.* (2000) [8], Patel *et al.* (2010) [12] and Bairagi (2014) [2] in fenugreek.

Table 1: Number of pods, seed per pod and seed yield as influenced by different nitrogen and phosphorus levels

Treatments	Number of pods plant ⁻¹	Seeds per pod	Seed yield (q ha ⁻¹)
Nitrogen levels			
N ₁ (40Kg ha ⁻¹)	13.14	9.01	8.24
N ₂ (60Kg ha ⁻¹)	15.37	10.19	10.72
N ₃ (80Kg ha ⁻¹)	27.33	13.64	13.27
N ₄ (100 Kg ha ⁻¹)	24.59	11.61	12.35
'F' test	Sig	Sig	Sig
S.E.(m)±	0.03	0.03	0.13
CD at 5%	0.10	0.10	0.37
Phosphorus levels			
P ₁ (20 Kgha ⁻¹)	17.24	10.61	10.15
P ₂ (40 Kgha ⁻¹)	17.87	11.08	11.32
P ₃ (60 Kgha ⁻¹)	18.47	11.65	11.96
'F test'	Sig.	Sig.	Sig.
S.E.(m)±	0.03	0.03	0.11
CD at 5%	0.08	0.09	0.32

Table 2: Number of pods, seed per pod and seed yield as influenced by interaction of different nitrogen and phosphorus levels

Treatments combination	Interaction Effect (N×P)		
	Number of pods plant ⁻¹	Seeds per pod	Seed yield (q ha ⁻¹)
N ₁ P ₁ (40 kg N + 20 kg P)	12.51	8.61	6.97
N ₁ P ₂ (40 kg N + 40 kg P)	13.16	8.95	8.38
N ₂ P ₃ (40 kg N + 60 kg P)	13.76	9.48	9.39
N ₂ P ₁ (60 kg N + 20 kg P)	14.74	9.74	9.81
N ₂ P ₂ (60 kg N + 40 kg P)	15.31	10.18	10.76
N ₂ P ₃ (60 kg N + 60 kg P)	16.07	10.66	11.58
N ₃ P ₁ (80 kg N + 20 kg P)	21.47	12.99	11.96
N ₃ P ₂ (80 kg N + 40 kg P)	22.20	13.68	13.50
N ₃ P ₃ (80 kg N + 60 kg P)	22.77	14.23	14.35
N ₄ P ₁ (100 kg N +20 kg P)	20.24	11.12	11.88
N ₄ P ₂ (100 kg N +40 kg P)	20.79	11.49	12.66
N ₄ P ₃ (100 kg N +60 kg P)	21.28	12.23	12.52
'F test'	NS	Sig.	Sig.
S.E.(m)±	0.05	0.05	0.22
CD at 5%	-	0.17	0.64

Test weight, germination % and protein content

Effect of nitrogen levels

A perusal of data on (Table 1) revealed that test weight, germination percent and protein content were significantly influenced by different levels of nitrogen. Significantly maximum test weight (23.55g) was recorded with 80 kg Nha⁻¹ and the minimum (17.72 g) with 40 kg Nha⁻¹. It was probably due to the fact that nitrogen have a vital role in the seed development and hence It has positively affected the seed weight up to the moderate dose of application. More available nitrogen filled the seeds and results the heavier seeds. The test weight or weight of 1000 seeds increases with increased level of nitrogen. Similar results were recorded by Patel *et al.* (2010) [12] and Kumar *et al.* (2015) [6]. The significantly maximum germination percent (94.33) was recorded with 100 kg Nha⁻¹ and the minimum germination percent (83.22) was observed with application of 40 kg Nha⁻¹. This might be due to the fact that nitrogen is primary nutrient required for protein synthesis and cell development so on germinating seeds it has positive effect. Increased nitrogen level enhanced germination %. These results confirms the finding of Sharma *et al.* (2000) [14] in fenugreek. Significantly maximum percent of Protein (23.68) was recorded with application of 80 kg Nha⁻¹. However, minimum percent of protein (20.26) was recorded with 40 kg Nha⁻¹. This was probably happened due to the nutrient can improve soil structure and root development, the optimum leaf area and better growth of plant with high nitrogen which led ultimately to better use of solar radiations and higher photosynthesis and finally highest

oil yield in seeds. results obtained in this study are in agreement with the study of Sunanda *et al.* (2014) [15] for fenugreek crop.

Effect of phosphorus

A perusal of data on (Table 1) revealed that test weight, germination percent and protein content were significantly influenced by different levels of phosphorus. Significantly maximum test weight (21.54g) was recorded with 60 kg Pha⁻¹. However, the minimum test weight(20.39g) was recorded with treatment 20 kg Pha⁻¹. Test weight was significantly influenced by different phosphorus levels. Phosphorus plays positive role in enhancement of food synthesis and its effective translocation towards the final photosyntates i.e. seed size and weight. Similar results were obtained by Nehara *et al.* (2002) [9] and Bairagi (2014) [2] in his research findind in fenugreek crop. The significantly maximum germination percentage (89.75) was recorded with treatment 60 kg Pha⁻¹, which was significantly superior over all other treatments. Whereas, minimum germination percentage (87.33) was recorded with 20 kg Pha⁻¹. An application of phosphorus also significantly increased the germination percent. This might be due to the fact that phosphorus are the primary nutrients required for protein synthesis and cell development. Phosphorus is called the energy forming nutrient as it increases carbohydrates of plant body and energy is required during seed germination so it provides energy to the germinating seeds. Phosphorus helped to establish seedling quickly and improved the quality of crop. Protein content was

significantly influenced by phosphorus levels. The maximum level of protein (22.79) was recorded with 60 kg Pha⁻¹, which was statistical. However, the minimum level of protein content (21.62) was recorded with 20 kg Nha⁻¹. Application of phosphorus has a positive effect on proper biomass production. Higher photosynthesis subsequently enhanced essential protein content. Protein plays important role in enzyme activity and metabolism. Similar result also obtained by Sunanda *et al.* (2014)^[15].

Interaction effect

Interaction effect due to different nitrogen and phosphorus levels on test weight was found to be significant. The maximum test weight (24.62) was recorded at treatment N₃P₃. However, minimum test weight (17.17) was recorded with

N₁P₁. similar results were obtained by Pareek and Gupta (1981)^[11], Rathore and Manohar (1989)^[13]. Interaction effect due to different nitrogen and phosphorus levels on germination % was found to be significant. The Significantly maximum germination % (95.33%) was recorded from treatment N₄P₃. Whereas the minimum (82.00%) was observed from treatment N₁P₁. Interaction effect due to different nitrogen and phosphorus levels on protein content was found to be significant. Significantly maximum content of protein (24.62%) was recorded with application of treatment N₃P₃. Whereas the minimum (19.69%) was observed from treatment N₁P₁. Results obtained in this study are in conformity with the study of Sunanda *et al.* (2014)^[15] on fenugreek.

Table 3: Quality parameters viz. test weight, germination per cent, protein content of seed as influenced by different nitrogen and phosphorus levels

Treatments	Test weight	Germination (%)	Protein content (%)
Nitrogen levels			
N ₁ (40Kg ha ⁻¹)	17.72	83.22	20.26
N ₂ (60Kg ha ⁻¹)	20.42	85.89	21.74
N ₃ (80Kg ha ⁻¹)	23.55	89.33	23.68
N ₄ (100 Kg ha ⁻¹)	22.36	94.33	23.29
'F test'	Sig.	Sig.	Sig.
S.E.(m)±	0.06	0.51	0.84
CD at 5%	0.19	1.51	2.45
Phosphorus levels			
P ₁ (20 Kg ha ⁻¹)	20.39	87.33	21.62
P ₂ (40 Kg ha ⁻¹)	21.12	87.50	22.32
P ₃ (60 Kg ha ⁻¹)	21.54	89.75	22.79
'F' test	Sig.	Sig.	NS
S.E.(m)±	0.05	0.45	0.72
CD at 5%	0.16	1.31	-

Table 4: Quality parameters viz. test weight, germination per cent and protein content as influenced by interaction of different nitrogen and phosphorus levels

Treatments combination	Interaction Effect (N×P)		
	Test weight	Germination (%)	Protein content (%)
N ₁ P ₁ (40 kg N + 20 kg P)	17.17	82.00	19.69
N ₁ P ₂ (40 kg N + 40 kg P)	17.73	82.67	20.29
N ₂ P ₃ (40 kg N + 60 kg P)	18.27	85.00	20.81
N ₂ P ₁ (60 kg N + 20 kg P)	19.29	86.67	21.25
N ₂ P ₂ (60 kg N + 40 kg P)	20.50	84.67	21.66
N ₂ P ₃ (60 kg N + 60 kg P)	21.47	86.33	22.30
N ₃ P ₁ (80 kg N + 20 kg P)	22.18	87.00	22.56
N ₃ P ₂ (80 kg N + 40 kg P)	23.85	88.67	23.85
N ₃ P ₃ (80 kg N + 60 kg P)	24.62	92.33	24.62
N ₄ P ₁ (100 kg N +20 kg P)	22.90	93.67	22.99
N ₄ P ₂ (100 kg N +40 kg P)	22.39	94.00	23.46
N ₄ P ₃ (100 kg N +60 kg P)	21.80	95.33	23.43
'F' test	Sig.	NS	NS
S.E.(m)±	0.11	0.89	1.45
CD at 5%	0.32	-	-

Conclusions

For yield and quality parameters the treatment combination (N₃P₃) T₈ i. e. 80 kg N + 20 kg P per hectare were found to be significantly superior. The significantly maximum pods per plant (22.77), seed yield per pod (14.23), seed yield per hectare (14.35q), test weight (24.62 g) and protein content (24.62%).

References

- Anonymous. NHB Data Base, 2014. <http://www.nhb.gov.in>

- Bairagi SK. Effect of different doses of phosphorus and row spacing on the yield and quality of fenugreek seed. The Asian J Hort. 2014; 9:338-341.
- Das PC. Vegetable crops of India. Hyderabad, Kalyani Publishers, 1992, 180-190.
- Chaudhary BR, Gupta AK, Parihar CM, Jat SL, Singh DK. Effect of intergrated nutrient management of fenugreek (*Trigonella foenum-graecum*) and its residual effect on fodder pearl millet. Indian J Agronomy. 2011; 56(3):185-195.

5. Jat BL. Effect of phosphorus, sulphur and biofertilizers on growth characters of fenugreek. (*Trigonella foenum-graecum* L.). Legume Res. 2001; 27(1):37-41.
6. Kumar R, Meena SS, Kakani RS, Mehta RS, Meena NK. Response of crop geometry, fertilizer levels and genotypes on productivity of fenugreek. International J Seed spice. 2015; 5(1):63-67.
7. Mehta RS, Anwer MM, Aishwath OP, Meena RS. Growth, yield and quality of fenugreek (*Trigonella foenum-graecum* L.) as influenced by nitrogen, phosphorus and bio-fertilizers. Indian J Hort. 2012; 69(1):94-9.
8. Mavai D, Lai S, Singh BA, Singh N, Mavai D. Response of fenugreek (*Trigonella foenum-graecum* L.) to seed rate, nitrogen and phosphorus fertilizer. Haryana J Hort. Sci. 2000; 29(3-4):244-246.
9. Nehara KC, Sharma US, Agarwal HR. Response of fenugreek (*Trigonella foenum-graecum* L.) to phosphorus and potassium. Annals of Biol. 2002; 18(1):43-45.
10. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. New Delhi, Indian Council of Agricultural Research, 1967, 215.
11. Pareek SK, Gupta R. Effect of fertilizer application on seed yield and diosgenin content in fenugreek. Indian J Agric. Sci. 1981; 50(10):746-749.
12. Patel BS, Patel SG, Patel SP, Amin AV. Intergrated Nutrient Management in fenugreek (*Trigonella foenum-graecum*) J. Spice and aromatic crops. 2010; 19(1&2):68-70.
13. Rathore PS, Manohar SS. Effect of date of sowing, levels of nitrogen and phosphorus on growth and yield of fenugreek. Madras Agric. J. 1989; 76(11):647-648.
14. Sharma SK. Response of nitrogen and spacing on fenugreek seed production. Hort. J. 2000; 13(2):39-42.
15. Sunanda BB, Shetty GR, Venkatesh J. Influence of integrated nutrient management on growth, yield and quality of Kasurimethi (*Trigonella corniculata* L.) under hill zone of Karnataka. International J Seed Spice. 2014; 4(2):62-67.
16. Sheoran RS, Sharma HC, Pannu PK. Efficiency of phosphorus fertilizer applied to fenugreek (*Trigonella foenum-graecum* L.) genotypes under different dates of sowing. HAU J Res. 1999; 29:101-107.